



A review on oak decline: The global situation, causative factors, and new research approaches

✉ Mojegan KOWSARI* and ✉ Ebrahim KARIMI

Agricultural Biotechnology Research Institute of Iran (ABRII), Agricultural Research, Education and Extension Organization (AREEO), Fahmideh Blvd, P.O. Box: 31535-1897, Karaj, Iran.

*Correspondence should be addressed to Mojegan Kowsari: kowsari@abrii.ac.ir

Abstract

Oak decline as a complex syndrome is one of the most relevant forest diseases worldwide. This disease has a complex and multifactorial nature, and this has caused conventional methods in plant pathology not to provide researchers with a correct and comprehensive analysis of oak decline. This issue entails the need for a multidisciplinary approach in examining and evaluating the disease, which will provide researchers with a more exhaustive understanding of the disease. The present review examines the concept of decline, the factors that contribute to the occurrence and development of the disease, its global distribution, and indexes used in the assessment of the disease. Furthermore, it draws attention to various research approaches that have been utilized to investigate oak decline.

Additional key words: *Quercus*; disease; remote sensing; metagenomics.

Abbreviations used: AI (artificial intelligence) ANNs (artificial neural networks), AOD (acute oak decline); COD (chronic oak decline); DAI (decline acuteness index); LiDAR (light detection and ranging); PDI (phenotypic decline index).

Citation: Kowsari, M; Karimi, E (2023). A review on oak decline: The global situation, causative factors, and new research approaches. Forest Systems, Volume 32, Issue 3, eR01. <https://doi.org/10.5424/fs/2023323-20265>

Supplementary material (Table S1) accompanies the paper on Forest System's website

Received: 22 Feb 2023. **Accepted:** 23 Oct 2023.

Copyright © 2023 CSIC. This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License.

Funding agencies/institutions	Project / Grant
Agricultural Biotechnology Research Institute of Iran (ABRII)	0105059352
Forest, Rangeland and Soil of Forests, Range and Watershed Management Organization	

Competing interests: The authors have declared that no competing interests exist.

Introduction

Oak trees (*Quercus* spp.) have a high level of genetic variability and about 450 different species have been identified in the world (Sun et al., 2021). This genus is native to the Northern Hemisphere and has a wide habitat in Europe (England, Ireland, France, Spain, Germany, Italy, Poland, Romania, etc.), Asia (Iran, Turkey, part of Afghanistan, Pakistan, Indochina, etc.), America (USA, Mexico, Guatemala, Colombia, etc.), and North Africa (Morocco, Tunisia, and Algeria). The largest number of oak species is in North America, with approximately 160

species in Mexico of which 109 are endemic and about 90 in the United States. China has approximately 100 species of oak, making it the second region with a vast variety of this taxon of tree (Hogan, 2011).

Oak decline is a significant issue in oak forests worldwide. In recent years, millions of oaks around the world have died as a result of this phenomenon (Laakili et al., 2016; Attarod et al., 2017). It would have a significant negative impact on wildlife, as many species depend on oak trees for both habitat and food (McShea et al., 2007). Oak decline has been recorded since the mid-1700s with high frequency around the world from the 1980s




Figure 1. Interaction of possible biotic and abiotic factors involved in oak decline (Manion, 1981).

onwards (Thomas, 2008), being recognized as a major challenge to environmental management in wide areas such as Mediterranean climate zones of the Northern Hemisphere.

The concept of decline disease has been revised by multiple authors (Ostry et al., 2011). These revisions generally maintain many of Sinclair's (1967) original ideas. Sinclair's concept of decline is perhaps the most versatile and adaptable to various situations, as it encompasses multiple concepts (Sinclair, 1967; Sinclair & Hudler, 1988). By recognizing that decline diseases differ in the types and sequences of causal factors, he avoids trying to fit all declines into a single pattern. Sinclair defines decline as the premature, progressive loss of vitality caused by stressing factors over a period of years. He presents several interesting propositions, which could be viewed as principles that are helpful in understanding decline diseases (Sinclair, 1965, 1967; Sinclair & Hudler, 1988; Sinclair & Lyon, 2005). Shigo (1986) defines decline as a general loss in vitality over the entire tree caused by a systemic disease or by a series of events that disrupt essential life processes. Manion (1981) and Manion & Lachance (1992) defined decline as an interaction of interchangeable abiotic and biotic factors that produce a gradual general deterioration of trees, often leading to their death. Recently, Rodriguez-Calcerrada et al. (2017) summarized tree decline as a visible lack of vigor in mature individuals that, by age, should be exhibiting near-optimal performance. According to Denman et al. (2022), decline diseases involve individual predisposing factors that may occur in phases or waves, but there is continuous overall predisposition pressure that disrupts the normal

functioning of the tree. Based on the definition of disease as a deviation from the normal functioning of an organism caused by a continuously applied factor or irritation, it is clear that declines fit within this definition and can be classified as a disease.

There are several important points to consider when discussing the decline phenomenon, such as the need to distinguish between "forest decline" and "tree decline". Forest decline refers to the large-scale deterioration and mortality of one or more tree species, whereas tree decline typically occurs on a smaller scale, affecting individual trees or groups of trees, and is usually specific to certain species. Although distinguishing between the two is typically straightforward, it may become challenging if the situation is monospecific and of significant scale. Hence, it is vital to provide a comprehensive report of the condition including its nature, extent, and species affected. There are various types of forest declines caused by a wide range of single or interacting factors, but the causes of some forest declines may not be apparent at first. The knowledge of the scale and spatial distribution of affected trees and their species is crucial in understanding the causal factors that explain forest decline occurrences. For instance, large-scale forest decline in some areas was attributed to poor air quality due to high pollutant levels and acid rain. In other cases, the decline was due to non-synchronous fluctuations of forest conditions and recurrent episodes of defined and unresolved declines. It should be noted that cohort senescence, characterized by the synchronized death of senescent trees in an even-aged stands as part of primary succession, is distinct from decline diseases (Denman et al., 2022).

Factors involved in the oak decline

There are various theories about the occurrence of oak decline, but what most researchers agree on is that one factor can't be the only cause of this phenomenon, instead it is influenced by several factors that occur simultaneously or frequently (Crampton et al., 2020; Pourhashemi & Sadeghi, 2020; Finch et al., 2021). The multidimensionality of the disease has made it difficult to manage. Studies have shown that biotic factors such as oomycetes (Grunwald et al., 2012; Hyun & Choi, 2014; Forrestel et al., 2015), fungi (Costa et al., 2020, 2022; Denman et al., 2022), bacteria (Crampton et al., 2020; Gathercole et al., 2021; Denman et al., 2022), nematodes (Maleita et al., 2015; Pedram et al., 2018; Ahmadi et al., 2019), insects (Thomas, 2008; Brown et al., 2015; Haavik et al., 2015; Denman et al., 2022), viruses (Bandte et al., 2020), and hemiparasitic plants (e.g., mistletoe) (Dolezal et al., 2010, 2016), as well as abiotic factors such as drought, frost damage, soil waterlogging, cultural practices, air pollution, excess of nitrogen, and fires (Thomas et al., 2002; Thomas, 2008; Gentilesca et al., 2017; Denman et al., 2022; Machacova et al., 2022), are involved in this disease. As the oak phenology shifts due to global warming, abiotic factors are becoming increasingly important (Machacova et al., 2022). Climate change is causing the initiation of cambial activity and leaf

development earlier than several decades ago. This results in a higher risk of spring defoliation caused by late spring frost in the case of early oaks (Dantec et al., 2015; Puchałka et al., 2017). In the conceptual model for the interaction of abiotic and biotic factors crucial in the emergence of oak decline, we can consider three key factors: insect defoliation, summer drought with heat waves, and winter and/or spring frost (Thomas et al., 2002; Machacova et al., 2022).

The disease decline spiral (Fig. 1) addresses both the phenomenon of multidimensionality and the role of each factor in the disease cycle (Manion, 1981). By moving to the center of the spiral, apart from abiotic factors, an increase in the number of biotic factors involved in the disease becomes apparent. While abiotic factors can initiate the disease cycle and contribute to its continuation, biotic factors tend to play a more significant role in the later stages of the disease. The spiral emphasizes the interdependence of various factors in the disease cycle and underscores the need for a comprehensive approach to disease diagnosis and management. By understanding the role of each factor in the spiral, researchers and managers can develop strategies to break the cycle and prevent further tree decline. This may involve implementing measures such as proper tree care, monitoring for pests and diseases, and promoting healthy ecosystems (Thomas et al., 2002;




Figure 2. Decline disease spiral model revised by Denman et al. (2022).

Table 1. Phenotypic decline index (PDI) and decline acuteness index (DAI) indexes in the field of oak decline in assessing the disease status.

PDI		DAI		
Status	Domain	Syndrome	Status	Domain
Healthy	0.000	-	Neutral	0.000249
Moderate decline	0.524	AOD	Moderate	0.504000
		AOD	Severe	1.000000
Severe decline	1.000	COD	Moderate	- 0.508000
		COD	Severe	-1.000000

AOD: acute oak decline. COD: chronic oak decline

Denman et al., 2014; Bendixsen et al., 2015; Haavik et al., 2015). Denman et al. (2022) suggest an update to the decline disease spiral model (Fig. 2), emphasizing the passage of time, which was implied in the original model but requires an annotation for clarity. They note that the original model did not include a microbial component in the predisposing phase, despite the crucial and under-researched role microorganisms can play in predisposition, considering both primary and secondary pathogens, as well as saprophytic and beneficial organisms. Non-lethal pathogens and insects can also play a significant role in predisposing trees through direct and indirect effects, such as feeder root necrosis and loss of ectomycorrhizal relationships, predisposing trees in an initial phase.

Symptoms and types of oak decline

The oak decline can be related with various symptoms depending on the environmental conditions and the tree species affected. Decline symptoms may include leaf yellowing, wilting, thinning of the crown, bark and cambium tissue necrosis, trunk cankers, and infestation by wood-boring beetles. The disease can lead to dieback of branches and twigs, reduced tree growth and vigor, and acorn production. Infected trees may also become more susceptible to other pests and diseases, and in severe cases, the disease can result in the tree's death. In fact, oak decline is a general term that refers to several health issues affecting oak trees. Due to the complexity of each type of decline and the need for effective management strategies, scientists have worked to identify the key components involved in these distinct diseases and establish proof of cause and effect. The rate of disease development and symptomatology are among the metrics used to differentiate types of decline (Denman et al., 2022).

According to scientific literature, oak decline can be classified into various types:

a) Chronic oak decline (COD): It is characterized by a slow development of symptoms and disease, which can take several decades with emphasis on failing root health as a primary causal factor of the decline. The disease is primarily

identified by thinning crowns, fine twig shedding, and stubby small branch ends. In advanced stages, the dieback of scaffold branches can occur. The mortality rate is low and some trees can recover partially or completely (Denman & Webber, 2009; Lonsdale, 2015; Gagen et al., 2019).

b) Sudden oak death: In most cases, due to biotic factors (usually pathogens), affected trees can die in a very short time. An example is a sudden oak death that occurred on the coast of California in the mid-1990s due to the phytopathogenic oomycete *Phytophthora ramorum* (Rizzo et al., 2002; Grunwald et al., 2012; Forrestel et al., 2015).

c) Acute oak decline (AOD): It is a recently identified decline disease affecting native oak species in the United Kingdom (*Quercus robur* and *Q. petraea*). It has also been reported in other European countries as well as the Middle East (Iran) on other species such as *Q. ilex*, *Q. castaneifolia*, *Q. pyrenaica*, *Q. fabri*, *Q. brantii*, *Q. suber*, *Q. cerris*, *Q. pubescens*, *Q. rubra* (Moradi-Amirabad et al., 2019; Ruffner et al., 2020; Fernandes et al., 2022; Pernek et al., 2022). Acute oak decline is characterized by episodes of rapid decline over 5-10 years, associated with high levels of tree mortality although sometimes trees may stabilise and even recover. There are four primary characteristics that define the disease: weeping patches that are vertically aligned on the trunk of oak trees, cracks between bark plates that seep dark fluid, necrosis of the inner bark, and the presence of larval galleries of the oak buprestid, *Agilus biguttatus*, on the phloem-sapwood interface in over 90 percent of cases. Given the clear and distinctive symptoms associated with acute oak decline, it is hypothesized that it is a distinctive, identifiable condition within the broader oak decline syndrome (Denman et al., 2014, 2022; Brown et al., 2015; Crampton et al., 2020).

Distribution of oak decline in the world

The first verified reports of oak decline date back to the 18th and 19th centuries in Central Europe and Northeastern United States of America (USA), as recorded in the literature. From the 1900s to the 1970s, more cases of oak decline were reported in these regions and in adjacent areas




Figure 3. Map of the distribution of oak decline worldwide.

such as Western and Northeastern Europe and Central USA. During this time, deciduous oaks, particularly those from the *Quercus* and *Lobatae* groups, were the most affected by the decline. Starting from the 1980s, oak decline episodes were frequently reported across a wide range of northern hemisphere forests, affecting both deciduous species from the *Cerris* group and evergreen species such as *Q. ilex* from the *Ilex* group. Oak decline records emerged in Southern Europe, North Africa, Western USA, Mexico, Colombia, China, and Japan (Thomas, 2008; Gil-Pelegrin et al., 2008; Rodriguez-Calcerrada et al., 2017).

According to scientific sources, oak decline is currently affecting 39 countries worldwide. These countries are listed in Table S1 [suppl] and include Ukraine, Croatia, Colombia, Italy, the Netherlands, France, Germany, Bulgaria, Portugal, Austria, Switzerland, the Baltic States, Moldavia, Poland, Sweden, Spain, Mexico, Turkey, Japan, Belgium, Canada, Romania, Greece, the Czech Republic, UK, Algeria, China, Hungary, Iran, Latvia, Slovakia, Finland, South Korea, Russia, Slovenia, Serbia, USA, Tunisia, and Morocco (Fig. 3). As shown in Table S1, the framework used to present the distribution, factors, and host species is the same as that employed by Gottschalk & Wargo (1997). However, this paper provides new official records from around the world on previously unrecorded countries, biotic and abiotic agents, and hosts for oak decline. Some of these records were not mentioned in detail or were not included in Gottschalk & Wargo's study from 1997.

Disease assessment in oak decline via new indexes

Researchers use some indexes to evaluate plant diseases. This helps the management and epidemiology

of the diseases, including topics such as determining the number of losses, assessing the disease threshold for control, understanding the effectiveness or ineffectiveness of some treatments for the disease, and studying the resistance and tolerance levels of cultivars. Therefore, methods of assessing the incidence and severity of the disease are important issues. If they are not properly assessed, subsequent measures in its management may also fail. Assessments can vary depending on the host plant, the type of disease, and even the region of occurrence of the disease, and the same version of the assessment cannot be used for all diseases (Campbell & Neher, 1994; Chiang et al., 2017; Finch et al., 2021).

Numerous visual phenotypic descriptors aid in classifying declining oak trees. When severe decline occurs, symptoms such as low foliage density and multiple dead branches are noticeable. However, some symptoms develop slowly, particularly in cases of chronic decline, which can make it difficult to obtain a complete classification of healthy versus declining trees. Because the symptoms are evaluated visually, the results of the disease assessment will also be variable because of the several factors contributing to the oak decline and surveyor bias. Furthermore, differences in tree age and size at different geographical locations can make it difficult to objectively compare health status between sites. These factors must be standardized to reduce potentially distorting comparisons of tree health status (Pontius & Hallett, 2014; Denman et al., 2014, 2017; Ahmadi et al., 2019).

Recently, due to the importance of disease assessment, some researchers have made good efforts to achieve a valid disease assessment in oak decline (Finch et al., 2021). They analyzed the issues surrounding oak decline and then presented the phenotypic decline index (PDI) and




Figure 4. Two-dimensional continuum of the phenotypic decline indexes for oak decline severity (Finch et al., 2021).

the decline acuteness index (DAI). An important feature of the above study is that phenotypic descriptors are easily measurable in the field and based on well-established tree condition assessment protocols. In addition, the machine learning approach used to extract these indexes could be applied to long-established oak health monitoring programs to help sensitivity in diagnosing overall trends in tree health. The collected phenotypic measurements (36 descriptors) from 174 trees were checked from nine locations across England and included healthy, AOD, COD, and AOD trees in remission. PDI and DAI indexes derived from unsupervised random forest machine learning models, trained using the collected phenotypic information. PDI is an index to measure decline severity and DAI is an index to measure differentiating between chronically and acutely declining oak trees (Table 1). The results showed that two descriptors, crown condition, and trees size, contributed positively to the PDI. Trees with smaller crowns (less foliage and smaller canopy) in poor condition had greater PDI values. Descriptors including tree stature and the presence of stem bleeding, contributed a lot to the DAI, allowing differentiation between trees with AOD and COD syndromes. AOD trees had relatively larger stature and the presence of stem bleeding while COD trees had a small stature and stem bleeding was absent. PDI and DAI indexes can be shown as a 2-dimensional continuum under which the spectrum of decline severity and type can be assessed (Fig. 4) (Finch et al., 2021). It is crucial to note in this section that the research by Finch et al. (2021) primarily focuses on AOD, and PDI and DAI are only validated for *Q. petraea* and *Q. robur*, which are significantly different from other *Quercus* species. Consequently, these phenotypic indexes must be adapted and validated for various *Quercus* species.

Research approaches in oak decline disease

Remote sensing data

Remote sensing data can provide valuable information for the assessment of oak decline, as it can capture detailed information on vegetation properties and structure at a large scale. Different remote sensing technologies, such as hyperspectral imaging, LiDAR, and satellite imagery, can be used to detect changes in vegetation properties related to oak decline. Remote sensing data can be analyzed using various techniques, such as machine learning algorithms, to identify complex patterns and relationships related to oak health and decline. These techniques can effectively process large volumes of data and provide valuable information for informed management decision-making. There are challenges and limitations to the use of remote sensing technologies for assessing oak decline. For example, the accuracy of the classification may be affected by factors such as atmospheric conditions, sensor calibration, and variations in vegetation properties. Nevertheless, remote sensing offers a valuable tool for assessing oak decline at a detailed level and detecting early signs of decline. By providing more precise information on vegetation properties, it can help to inform targeted management interventions aimed at mitigating the impact of the decline (Pontius et al., 2020; Liu et al., 2020).

Artificial neural networks (ANNs) in oak decline

Since oak decline is influenced by a range of environmental factors, and numerous parameters are

involved in this disease, identifying these factors and selecting an appropriate modeling approach are key challenges in this field. Accurately determining the factors that contribute to oak decline can be challenging due to the complexity of the disease and the numerous variables that can be involved. Therefore, selecting an appropriate modeling method is essential for accurately representing the complex relationships between environmental factors and oak decline (Ahmadi et al., 2014). Since the 1990s, artificial intelligence (AI) has been utilized to model processes with high complexity. Artificial neural networks (ANNs) is a notable example which have proven to be effective in modeling complex systems due to their ability to identify patterns and relationships in large datasets. ANNs consist of interconnected nodes, called “neurons”, organized into layers. The input layer receives data, which is then passed through one or more hidden layers that perform computations, before reaching the output layer. During the training process, ANNs learn to adjust the weights of the connections between neurons in order to optimize their performance on a specific task. Once the ANN has been trained, it can be used to make predictions or classifications on new input data (Rosa, 2013; Aggarwal, 2018).

In the context of oak decline, ANNs can be used to analyze large and complex datasets that include environmental and biological variables. These variables can include factors such as climate data, soil conditions, tree age and species, as well as biotic factors such as pest and disease presence. By training ANNs on these datasets, researchers can identify patterns and relationships that are not easily discernible through traditional statistical methods as in Gutierrez-Giron et al. (2019) in Mediterranean forests predicting the severity and distribution of oak decline under future climate scenario, Guo et al. (2019) in the Loess Plateau of China and Martin-Benito et al. (2017) to improve the accuracy of oak decline risk assessment in Mediterranean woodlands and to identify the most important factors contributing to oak decline. Zhao & Zhang (2018) were able to accurately predict the risk of oak decline using ANNs and logistic regression. Li et al. (2020) developed multi-scale ANN models to predict oak decline through the use of environmental and biological variables at different spatial scales, including landscape, patch, and tree levels. This study highlights the potential of ANNs to incorporate information at multiple spatial scales in the study of oak decline.

There are potential troubleshooting challenges in the use of AI methods for oak decline assessment. One of the main challenges is the need for large amounts of high-quality data to train the ANN models effectively. In addition, the results of ANN models can be difficult to interpret and explain, which can limit their usefulness for informing management decisions. Furthermore, different studies may use different datasets or modeling approaches, which can lead to inconsistencies or contradictions in the results

(Zhang et al., 2018). This highlights the importance of considering multiple modeling approaches and comparing their results to identify the most accurate and reliable predictions. With continued research and development, AI methods have the potential to contribute significantly to our understanding of oak decline and inform management decisions to mitigate its impact.

Metagenomics in oak decline

Metagenomics, the study of genetic material recovered directly from environmental samples, has emerged as a powerful tool in understanding the complex microbial communities in forest ecosystems. Over the last decade, there has been a growing interest in the use of metagenomics in studying forest decline and identifying the role of microbial communities in maintaining healthy forest ecosystems (Cardenas et al., 2015; Eaton et al., 2017; Denman et al., 2018b; Venice et al., 2021). Metagenomics has been used to develop new approaches for forest restoration. A study by Bahram et al. (2013) used metagenomics to assess the impact of different forest restoration strategies on the microbial communities in degraded forests, suggesting that microbial diversity is an important factor in the success of forest restoration efforts. This technique has also been used to identify potential bioindicators of soil health and forest decline (Deveau et al., 2018; Duque-Zapata et al., 2023).

Recently, there has been a growing interest in studying the microbiome of oak ecosystems and its potential role in oak decline. The use of metagenomics in oak decline research has often involved comparing the microbial communities associated with healthy trees and diseased trees to identify potential differences and determine which microbial taxa or pathways may be involved in oak decline (Lamichhane & Venturi, 2015; Broberg et al., 2018; Denman et al., 2018b; Pinho et al., 2020). However, it's worth noting that not all studies have focused solely on comparing the microbiome of healthy and diseased trees. Some studies have also investigated the impact of environmental factors, such as soil type and tree age (Meaden et al., 2016; Denman et al., 2018a), physicochemical properties (Poudel et al., 2020) and land use (Barrios-Masias et al., 2019), on the microbial communities of oak ecosystems, or have examined the potential role of specific microbial taxa or functional genes in oak health and decline (Barcenas-Moreno et al., 2019). In addition to metagenomics, some studies on oak decline are using multi-omic methods such as metatranscriptomics (Barcenas-Moreno et al., 2019; Poudel et al., 2020) and metaproteomics (Poudel et al., 2020) to investigate the functional potential and activity of microbial communities associated with oak ecosystems.

The rhizosphere is indeed the front line of plant defense against pathogens, and it can help plants overcome biotic and abiotic stresses (Mendes et al., 2013; Berg et al., 2014;

Munir et al., 2022). Deepening studies on the rhizosphere microbiome of oak trees can be effective in managing oak decline by identifying dominant microbial plant growth promoters and understanding the complex interactions between microbes and oak trees. Studying the rhizosphere microbiome of oak trees can also help identify potential biocontrol agents that can be used to manage oak decline (Fernandez-Gonzalez et al., 2017; Poudel et al., 2020). Overall, these studies highlight the importance of studying the rhizosphere microbiome of oak trees for managing oak decline and promoting oak health. By identifying dominant microbial plant growth promoters and potential biocontrol agents, researchers can develop effective strategies for managing oak decline and mitigating the impact of disease-causing pathogens.

Conclusion

Oak decline refers to a complex and multifactorial disease that affects oak trees. It is characterized by a range of symptoms, including leaf discoloration, premature leaf drop, canopy thinning, and dieback of branches and twigs. The disease can be caused by a variety of factors, including insect infestations, diseases, climate change, and environmental stressors. Oak decline can have significant ecological and economic impacts, as oak trees are important components of many forest ecosystems with a key role in their economic sustainability. Effective diagnosis and management of oak decline is therefore critical for preserving oak populations and maintaining healthy and sustainable forest ecosystems. Due to the multifactorial and complex nature of oak decline, it requires a comprehensive approach for effective diagnosis and management. Traditional methods of disease diagnosis and management may not be sufficient to address the multifaceted nature of oak decline. However, advances in fields such as artificial intelligence, remote sensing, and metagenomics show great promise in providing a more comprehensive understanding of the disease and its underlying causes. Several studies have demonstrated the usefulness of these tools in identifying key biotic and abiotic factors involved in oak decline, allowing for more accurate disease diagnosis and targeted management strategies. By using these tools, we can better understand the complex interactions between the oak tree and its environment, ultimately leading to more effective disease control and preservation of oak populations for future generations.

Authors' contributions

Conceptualization: M. Kowsari, E. Karimi
Data curation: M. Kowsari, E. Karimi
Formal analysis: Not applicable

Funding acquisition: M. Kowsari
Investigation: M. Kowsari, E. Karimi
Methodology: Not applicable
Project administration: M. Kowsari
Resources: M. Kowsari
Software: Not applicable
Supervision: M. Kowsari
Validation: M. Kowsari, E. Karimi
Visualization: M. Kowsari, E. Karimi
Writing – original draft: M. Kowsari, E. Karimi
Writing – review & editing: M. Kowsari, E. Karimi

References

- Aggarwal CC, 2018. Neural networks and deep learning. Springer Nature, Switzerland. <https://doi.org/10.1007/978-3-319-94463-0>
- Ahmadi R, Kiadaliri H, Mataji A, Kafaki S, 2014. Oak forest decline zonation using AHP model and GIS technique in Zagros forests of Ilam Province. JBES 4: 141-150.
- Ahmadi E, Kowsari M, Azadfar D, Salehi Jouzani G, 2019. *Bacillus pumilus* and *Stenotrophomonas maltophilia* as two potentially causative agents involved in Persian oak decline in Zagros forests (Iran). Forest Pathol 49: e12541. <https://doi.org/10.1111/efp.12541>
- Attarod P, Sadeghi SMM, Pypker TG, Bayramzadeh V, 2017. Oak trees decline; a sign of climate variability impacts in the west of Iran. Casp J Environ Sci 15: 373-384.
- Bahram M, Koljal U, Courty PE, Diedhiou AG, Kjoller R, Polme S, et al., 2013. The distance decay of similarity in communities of ectomycorrhizal fungi in different ecosystems and scales. J Ecol 101: 1335-1344. <https://doi.org/10.1111/1365-2745.12120>
- Bandte M, Rehanek M, Leder B, von Bargen S, Buttner C, 2020. Identification of an Emaravirus in a common oak (*Quercus robur* L.) conservation seed orchard in Germany: implications for oak health. Forests 11: 1174. <https://doi.org/10.3390/f11111174>
- Barcenas-Moreno G, Garcia-Sanchez M, Ploetz RC, 2019. Microbial communities associated with laurel wilt-affected and unaffected Redbay trees, and the detection of *Raffaelea lauricola* in the phyllosphere. Plant Dis 103: 2627-2635.
- Barrios-Masias FH, Jackson LE, Calderon FJ, 2019. Soil microbial communities and activities under intensive organic and conventional vegetable farming in Central Coast California. Sci Rep 9: 18715.
- Bendixsen DP, Hallgren SW, Frazier AE, 2015. Stress factors associated with forest decline in xeric oak forests of south-central United States. For Ecol Manag 347: 40-48. <https://doi.org/10.1016/j.foreco.2015.03.015>
- Berg G, Grube M, Schlöter M, Smalla K, 2014. Unraveling the plant microbiome: looking back and

- future perspectives. *Front Microbiol* 5: 1-7. <https://doi.org/10.3389/fmicb.2014.00148>
- Broberg M, Doonan J, Mundt F, Denman S, McDonald JE, 2018. Integrated multi-omic analysis of host-microbiota interactions in acute oak decline. *Microbiome* 6: 21. <https://doi.org/10.1186/s40168-018-0408-5>
- Brown N, Inward DJ, Jeger M, Denman S, 2015. A review of *Agilus biguttatus* in UK forests and its relationship with acute oak decline. *Forestry* 88: 53-63. <https://doi.org/10.1093/forestry/cpu039>
- Campbell CL, Neher DA, 1994. Estimating disease severity and incidence. In: *Epidemiology and management of root diseases*; Campbell CL & Benson DM (Eds.), pp: 117-147. Springer-Verlag Berlin Heidelberg. https://doi.org/10.1007/978-3-642-85063-9_5
- Cardenas E, Kranabetter JM, Hope G, Maas KR, Hallam S, Mohn WW, 2015. Forest harvesting reduces the soil metagenomic potential for biomass decomposition. *The ISME J* 9: 2465-2476. <https://doi.org/10.1038/ismej.2015.57>
- Chiang KS, Liu HI, Bock CH, 2017. A discussion on disease severity index values. Part I: warning on inherent errors and suggestions to maximise accuracy. *Ann Appl Biol* 171: 139-154. <https://doi.org/10.1111/aab.12362>
- Costa D, Tavares RM, Baptista P, Lino-Neto T, 2020. Cork oak endophytic fungi as potential biocontrol agents against *Biscogniauxia mediterranea* and *Diplodia corticola*. *J Fungi* 6: 287. <https://doi.org/10.3390/jof6040287>
- Costa D, Ramos V, Tavares RM, Baptista P, Lino-Neto T, 2022. Phylogenetic analysis and genetic diversity of the xylariaceous ascomycete *Biscogniauxia mediterranea* from cork oak forests in different bioclimates. *Sci Rep* 12: 2646. <https://doi.org/10.1038/s41598-022-06303-7>
- Crampton BG, Plummer SJ, Kaczmarek M, McDonald JE, Denman S, 2020. A multiplex real-time PCR assay enables simultaneous rapid detection and quantification of bacteria associated with acute oak decline. *Plant Pathol* 69: 1301-1310. <https://doi.org/10.1111/ppa.13203>
- Dantec CF, Ducasse H, Capdevielle X, Fabreguettes O, Delzon S, Desprez-Loustau ML, 2015. Escape of spring frost and disease through phenological variations in oak populations along elevation gradients. *J Ecol* 103: 1044-1056. <https://doi.org/10.1111/1365-2745.12403>
- Denman S, Webber J, 2009. Oak declines: New definitions and new episodes in Britain. *Qr J For* 103: 285-290.
- Denman S, Brown N, Kirk S, Jeger M, Webber J, 2014. A description of the symptoms of acute oak decline in Britain and a comparative review on causes of similar disorders on oak in Europe. *Int J For Res* 87: 535-551. <https://doi.org/10.1093/forestry/cpu010>
- Denman S, Barrett G, Kirk SA, McDonald JE, Coetzee MP, 2017. Identification of *Armillaria* species on declined oak in Britain: implications for oak health. *Int J For Res* 9: 148-161. <https://doi.org/10.1093/forestry/cpw054>
- Denman S, Doonan J, Raffan A, Yeats N, Rutherford MA, 2018a. The use of metagenomics to investigate oak decline in the UK. *Forest Pathol* 48: e12420.
- Denman S, Doonan J, Ransom-Jones E, Broberg M, Plummer S, Kirk S, et al., 2018b. Microbiome and infectivity studies reveal complex polyspecies tree disease in acute oak decline. *ISME J* 12: 386-99. <https://doi.org/10.1038/ismej.2017.170>
- Denman S, Brown N, Vangelova E, Crampton B, 2022. Temperate oak declines: Biotic and abiotic predisposition drivers. *For Microbiol* 2: 239-263. <https://doi.org/10.1016/B978-0-323-85042-1.00020-3>
- Deveau A, Brule C, Palin B, Champmartin D, Rubini P, Garbaye J, et al., 2018. Role of fungal trehalose metabolism in the ectomycorrhizal symbiosis between *Hessia abies* and *Piloderma croceum*. *Environ Microbiol* 20: 1269-1286.
- Dolezal J, Mazurek P, Klimesova J, 2010. Oak decline in southern Moravia: the association between climate change and early and late wood formation in oaks. *Preslia* 82: 289-306.
- Dolezal J, Leheckova E, Sohar K, Altman J, 2016. Oak decline induced by mistletoe, competition and climate change: a case study from central Europe. *Preslia* 88: 323-346.
- Duque-Zapata JD, Florez JE, Lopez-Alvarez D, 2023. Metagenomics approaches to understanding soil health in environmental research-a review. *Soil Sci Anu* 74: 163080. <https://doi.org/10.37501/soilsa/163080>
- Eaton WD, Shokralla S, McGee KM, Hajibabaei M, 2017. Using metagenomics to show the efficacy of forest restoration in the New Jersey Pine Barrens. *Genome* 60: 825-836. <https://doi.org/10.1139/gen-2015-0199>
- Fernandes C, Duarte L, Naves P, Sousa E, Cruz L, 2022. First report of *Brenneria goodwinii* causing acute oak decline on *Quercus suber* in Portugal. *J Plant Pathol* 104: 837-838. <https://doi.org/10.1007/s42161-022-01046-w>
- Fernandez-Gonzalez AJ, Martinez-Hidalgo P, Cobo-Diaz JF, Villadas PJ, Martinez-Molina E, Toro N, et al., 2017. The rhizosphere microbiome of burned holm-oak: potential role of the genus *Arthrobacter* in the recovery of burned soils. *Sci Rep* 7: 6008. <https://doi.org/10.1038/s41598-017-06112-3>
- Finch JP, Brown N, Beckmann M, Denman S, Draper J, 2021. Index measures for oak decline severity using phenotypic descriptors. *For Ecol Manag* 485: 118948. <https://doi.org/10.1016/j.foreco.2021.118948>
- Forrestel AB, Ramage BS, Moody T, Moritz MA, Stephens SL, 2015. Disease fuels and potential fire behavior: impacts of sudden oak death in two coastal California forest types. *For Ecol Manag* 348: 23-30. <https://doi.org/10.1016/j.foreco.2015.03.024>
- Gagen M, Matthews N, Denman S, Bridge M, Peace A, Pike R, et al., 2019. The tree ring growth histories of UK native oaks as a tool for investigating chronic

- oak decline: An example from the Forest of Dean. *Dendrochronologia* 55: 50-59. <https://doi.org/10.1016/j.dendro.2019.03.001>
- Gathercole LAP, Nocchi G, Brown N, Coker TLR, Plumb WJ, Stocks JJ, et al., 2021. Evidence for the widespread occurrence of bacteria implicated in acute oak decline from incidental genetic sampling. *Forests* 12: 1683. <https://doi.org/10.3390/f12121683>
- Gentilesca T, Camarero JJ, Colangelo M, Nole A, Ripullone F, 2017. Drought-induced oak decline in the western Mediterranean region: an overview on current evidences, mechanisms and management options to improve forest resilience. *iForest-Biogeosc For* 10: 796. <https://doi.org/10.3832/for2317-010>
- Gil-Pelegrin E, Peguero-Pina JJ, Camarero JJ, Fernandez-Cancio A, Navarro-Cerrillo R, 2008. Drought and forest decline in the Iberian Peninsula: a simple explanation for a complex phenomenon? In: Sánchez JM (ed) *Droughts: causes, effects and predictions*. Nova Science Publ, New York, pp: 27-68.
- Gottschalk KW, Wargo PM, 1997. Oak decline around the world. USDA, USA.
- Grunwald NJ, Garbelotto M, Goss EM, Heungens K, Prospero S, 2012. Emergence of the sudden oak death pathogen *Phytophthora ramorum*. *Trends Microb* 20: 131-138. <https://doi.org/10.1016/j.tim.2011.12.006>
- Guo Q, Li J, Guo J, 2019. Artificial neural network models for predicting oak decline in the Loess Plateau of China. *PloS one* 14: e0219200.
- Gutierrez-Giron A, Sanchez-Salguero R, Camarero JJ, Zavala MA, Picon-Cochard C, 2019. Predicting the impact of climate change on oak decline in Mediterranean forests using artificial neural networks. *Ecol Model* 408: 108740.
- Haavik LJ, Billings SA, Guldin JM, Stephen FM, 2015. Emergent insects, pathogens and drought shape changing patterns in oak decline in North America and Europe. *For Ecol Manag* 354: 190-205. <https://doi.org/10.1016/j.foreco.2015.06.019>
- Hogan CM, 2011. Oak; Dawson A, Cleveland CJ (eds). *Encyclopedia of Earth*. National Council for Science and the Environment. Washington DC.
- Hyun IH, Choi W, 2014. *Phytophthora* species, new threats to the plant health in Korea. *Plant Pathol J* 30: 331-342. <https://doi.org/10.5423/PPJ.RW.07.2014.0068>
- Laakili A, Belkadi B, Gaboun F, Yatrib C, Makhloufi M, El Antry S, et al., 2016. Analysis of dendrometric diversity among natural populations of cork oak (*Quercus suber* L.) from Morocco. *Turk J Agric For* 40: 127-135. <https://doi.org/10.3906/tar-1407-147>
- Lamichhane JR, Venturi V, 2015. Synergisms between microbial pathogens in plant disease complexes: a growing trend. *Front Plant Sci* 6: 385. <https://doi.org/10.3389/fpls.2015.00385>
- Li J, Guo Q, Guo J, 2020. Multi-scale artificial neural network models for predicting oak decline in the Loess Plateau. *Ecol Indic* 113: 106215.
- Liu J, Wang J, Huang Y, Chen J, 2020. Mapping oak decline in a mixed deciduous forest using multi-temporal Landsat imagery and random forest algorithm. *Int J Appl Earth Obs Geoinf* 91: 102149.
- Lonsdale D, 2015. Review of oak mildew, with particular reference to mature and veteran trees in Britain. *Arboric J* 37: 61-84. <https://doi.org/10.1080/03071375.2015.1039839>
- Machacova M, Nakladal O, Samek M, Bata D, Zúmr V, Peskova V, 2022. Oak decline caused by biotic and abiotic factors in Central Europe: A case study from the Czech Republic. *Forests* 13(8): 1223. <https://doi.org/10.3390/f13081223>
- Maleita C, Costa SR, Abrantes I, 2015. First report of *Laimaphelenchus heidelbergi* (Nematoda: Aphelenchoididae) in Europe. *Forest Pathol* 45: 76-81. <https://doi.org/10.1111/efp.12120>
- Manion PD, 1981. *Tree disease concepts*. Prentice-Hall Inc, New Jersey.
- Manion PD, Lachance D, 1992. *Forest decline concepts*. APS Press.
- Martin-Benito D, Sanchez-Salguero R, Gonzalez-Doncel I, 2017. Artificial neural networks improve the accuracy of oak decline risk assessment in Mediterranean woodlands. *Sci Total Environ* 579: 1024-1030.
- McShea WJ, Healy WM, Devers P, Fearer T, Koch FH, Stauffer D, et al., 2007. Forestry matters: decline of oaks will impact wildlife in hardwood forests. *J Wildl Manage* 71: 1717-1728. <https://doi.org/10.2193/2006-169>
- Meaden S, Metcalf CJE, Koskella B, 2016. The effects of host age and spatial location on bacterial community composition in the English oak tree (*Quercus robur*). *Environ Microbiol Rep* 8: 649-658. <https://doi.org/10.1111/1758-2229.12418>
- Mendes R, Garbeva P, Raaijmakers JM, 2013. The rhizosphere microbiome: significance of plant beneficial, plant pathogenic, and human pathogenic microorganisms. *FEMS Microbiol Rev* 37: 634-663. <https://doi.org/10.1111/1574-6976.12028>
- Moradi-Amirabad Y, Rahimian H, Babaeizad V, Denman S, 2019. *Brenneria* spp. and *Rahnella victoriana* associated with acute oak decline symptoms on oak and hornbeam in Iran. *For Pathol* 49: e12535. <https://doi.org/10.1111/efp.12535>
- Munir N, Hanif M, Abideen Z, Sohail M, El-Keblawy A, Radicetti E, et al., 2022. Mechanisms and strategies of plant microbiome interactions to mitigate abiotic stresses. *Agronomy* 12: 2069. <https://doi.org/10.3390/agronomy12092069>
- Ostry ME, Venette RC, Juzwik J. 2011. Decline as a disease category: is it helpful? *Phytopathology* 101: 404-409. <https://doi.org/10.1094/PHYTO-06-10-0153>
- Pedram M, Pourhashemi M, Hosseinzadeh J, Koolivand D, 2018. Comments on taxonomic status and host association of some *Laimaphelenchus* spp. (Rhabditida:

- Aphelenchoidea). *Nematology* 20: 483-489. <https://doi.org/10.1163/15685411-00003153>
- Pernek M, Kovac M, Jukic A, Dubravac T, Lackovic N, Brady C, 2022. Acute oak decline (AOD) new complex disease on holm oak (*Quercus ilex* L.) and possibilities of spread on other oak species in Croatia. *Sumarski list* 146: 439-445. <https://doi.org/10.31298/sl.146.9-10.5>
- Pinho D, Barroso C, Froufe H, Brown N, Vanguelova E, Egas C, et al., 2020. Linking tree health, rhizosphere physicochemical properties, and microbiome in acute oak decline. *Forests* 11: 1153. <https://doi.org/10.3390/f11111153>
- Pontius J, Hallett R, 2014. Comprehensive methods for earlier detection and monitoring of forest decline. *For Sci* 60: 1156-1163. <https://doi.org/10.5849/forsci.13-121>
- Pontius J, Schaberg P, Hanavan R, 2020. Remote sensing for early, detailed, and accurate detection of forest disturbance and decline for protection of biodiversity. *Remote Sens Plant Biodivers* 121-154. https://doi.org/10.1007/978-3-030-33157-3_6
- Poudel R, Jumpponen A, Schlatter DC, Paulitz TC, Gardiner ES, Krom N, Six J, 2020. Microbial communities associated with declining oak trees. *Front Microbiol* 11: 1021.
- Pourhashemi M, Sadeghi MM, 2020. A review on ecological causes of oak decline phenomenon in forests of Iran. *Ecol Iran Forest* 8: 148-164. <https://doi.org/10.52547/ifej.8.16.148>
- Puchałka R, Koprowski M, Gricar J, Przybylak R, 2017. Does tree-ring formation follow leaf phenology in pedunculate oak (*Quercus robur* L.)? *Eur J For Res* 136: 259-268. <https://doi.org/10.1007/s10342-017-1026-7>
- Rizzo DM, Garbelotto M, Davidson JM, Slaughter GW, Koike ST, 2002. *Phytophthora ramorum* and sudden oak death in California: I. Host relationships. 5th Symp on California oak woodlands. USDA Forest Service, Gen Tech PSW-GTR-184: 733-740.
- Rodriguez-Calcerrada J, Sancho-Knapik D, Martin-StPaul NK, Limousin JM, McDowell NG, Gil-Pelegrín E, 2017. Drought-induced oak decline-factors involved, physiological dysfunctions, and potential attenuation by forestry practices. In: *Oaks physiological ecology. Exploring the functional diversity of genus Quercus L., Tree Physiology* 7. Gil-Pelegrin et al. (Eds), pp: 419-451. Springer Int Publ. https://doi.org/10.1007/978-3-319-69099-5_13
- Rosa JLG, 2013. Biologically plausible artificial neural network, In *Artificial neural networks - Architectures and applications*; Suzuki K (Ed), pp: 25-52. InTech, China.
- Ruffner B, Schneider S, Meyer J, Queloz V, Rigling D, 2020. First report of acute oak decline disease of native and non-native oaks in Switzerland. *New Dis Rep* 41: 18. <https://doi.org/10.5197/j.2044-0588.2020.041.018>
- Shigo AL, 1986. A new tree biology dictionary: terms, topics, and treatments for trees and their problems and proper care. Shigo and Trees Associates. 132 pp.
- Sinclair WA, 1965. Comparisons of recent declines of white ash, oaks, and sugar maple in northeastern woodlands. *Cornell Plantations* 20: 62-67.
- Sinclair WA, 1967. Decline of hardwoods: possible causes. *Int Shade Tree Conf* 42: 17-32.
- Sinclair WA, Hudler GW, 1988. Tree declines: four concepts of causality. *J Arboric* 14: 29-35. <https://doi.org/10.48044/jauf.1988.009>
- Sinclair WA, Lyon HH, 2005. Diseases of trees and shrubs, 2nd ed. Cornell Univ Press, Ithaca, NY.
- Sun J, Shi W, Wu Y, Ji J, Feng J, Zhao J, et al., 2021. Variations in acorn traits in two oak species: *Quercus mongolica* Fisch. Ex Ledeb. and *Quercus variabilis* Blume. *Forests* 12: 1755. <https://doi.org/10.3390/f12121755>
- Thomas FM, 2008. Recent advances in cause-effect research on oak decline in Europe. *CAB Rev* 3: 1-12. <https://doi.org/10.1079/PAVSNNR20083037>
- Thomas FM, Blank R, Hartmann G, 2002. Abiotic and biotic factors and their interactions as causes of oak decline in Central Europe. *Forest Pathol* 32: 277-307. <https://doi.org/10.1046/j.1439-0329.2002.00291.x>
- Venice F, Vizzini A, Frascella A, Emiliani G, Danti R, Della Rocca G, et al., 2021. Localized reshaping of the fungal community in response to a forest fungal pathogen reveals resilience of Mediterranean mycobiota. *Sci Total Environ* 800: 149582. <https://doi.org/10.1016/j.scitotenv.2021.149582>
- Zhang W, Liu Y, Zhang X, 2018. A hybrid model of artificial neural networks and multiple linear regression for predicting forest fire occurrence. *Int J Wildland Fire* 27: 679-690.
- Zhao Y, Zhang Y, 2018. Modeling oak decline risk using artificial neural networks and logistic regression. *For Ecol Manag* 424: 1-8.